

Electric properties of the baryon anti-decuplet in the SU(3) chiral quark-soliton model

T. Ledwig,^{1,*} H.-Ch. Kim,^{2,†} A. J. Silva,^{3,4,‡} and K. Goeke^{1,§}

¹*Institut für Theoretische Physik II,*

Ruhr-Universität Bochum, D-44780 Bochum, Germany

²*Department of Physics, and Nuclear Physics & Radiation Technology Institute (NuRI),*

Pusan National University, 609-735 Busan, Republic of Korea

³*Departamento de Física and Centro de Física Computacional,*

Universidade de Coimbra, P-3000 Coimbra, Portugal

⁴*Faculdade de Engenharia da Universidade do Porto,*

R. Dr. Roberto Frias s/n, P-4200-465 Porto, Portugal

(Dated: March 2006)

Abstract

We investigate the electric form factors and charge radii of the pentaquark baryons within the framework of the chiral quark-soliton model. We consider the rotational $1/N_c$ and linear m_s corrections, assuming isospin symmetry and employing the symmetry-conserving quantization. The flavor-decomposed charge densities of the Θ^+ are presented. The electric form factors and charge radii of the charged pentaquark baryons turn out to be very similar to those of the corresponding octet baryons. The charge radii of the neutral pentaquark baryons are obtained to be very tiny and positive. The strange electric form factor of the pentaquark proton is shown to be larger than the corresponding one of the proton by around 20 %. We also present the charge radii of the baryon decuplet for comparison.

PACS numbers: 12.39.Fe, 13.40.Em, 12.40.-y, 14.20.Dh

Keywords: Pentaquark baryons, electric form factors, charge radii, chiral quark-soliton model

*Electronic address: Tim.Ledwig@tp2.rub.de

†Electronic address: hchkim@pusan.ac.kr

‡Electronic address: ajose@teor.fis.uc.pt

§Electronic address: Klaus.Goeke@tp2.rub.de

1. The pentaquark baryons have been a hot issue, since Diakonov *et al.* predicted the mass and width of the pentaquark baryon Θ^+ [1] with strange quark number $S = +1$ and valence quark structure $uudd\bar{s}$. An earlier estimate of the mass in the soliton approach of the Skyrme model was given by Praszalowicz [2]. A great deal of experimental and theoretical works have been already piled up to confirm its existence and to understand its properties (see, for example, a recent review [3] for the experimental results). Many experiments have announced the existence of the Θ^+ after the first observation by the LEPS collaboration [4], while the Θ^+ has not been seen in almost all high-energy experiments. Moreover, exotic Ξ_{T0} states were observed by the NA49 experiment at CERN [5], though its existence is still under debate.

A very recent CLAS experiment, a dedicated experiment to search for the Θ^+ , has announced null results of finding the Θ^+ in $\gamma p \rightarrow \bar{K}^0 \Theta^+$ [6]. Though this experiment is high statistics measurement, it is too early to conclude the absence of the Θ^+ , because the previous positive evidences appeared mostly in the reactions from the neutron and the kinematic regions of each experiment are different. Moreover, several new experiments searching for the Θ^+ are in progress [7, 8]. In the present obscure status for the Θ^+ , more efforts should be made for understanding the Θ^+ theoretically as well as experimentally.

In addition, a recent GRAAL experiment [9] announced a new nucleon-like resonance with a seemingly narrow decay width ~ 10 MeV and a mass ~ 1675 MeV in the η -photoproduction from the neutron target. This new nucleon-like resonance, $N^*(1675)$, may be regarded as a non-strange pentaquark because of its narrow decay width which is known to be one of the characteristics of typical pentaquark baryons, though one should not exclude a possibility that it might be one of the known πN resonances (possibly, D_{15}) [10]. This GRAAL data is consistent with the results for the transition magnetic moments in the chiral quark-soliton model (χ QSM) [11] as well as the partial-wave analysis for the non-strange pentaquark baryons [12]. Moreover, a recent theoretical calculation of the $\gamma N \rightarrow \eta N$ reaction [13] describes qualitatively well the GRAAL data, based on the values of the magnetic transition moments in Refs. [11, 12], which implies that the N^* seen in the GRAAL experiment could be favorably identified as one of the pentaquark baryons.

Being motivated by these new experimental findings, a great deal of theoretical investigations have been performed, which are summarized in recent reviews [14, 15, 16]. Much of these theoretical works has concentrated on understanding the structure of these exotic pentaquark baryons. While the mass of the Θ^+ is experimentally known, other properties such as its spin and parity are undetermined. Moreover, the electromagnetic properties like charge radii and magnetic moments, and various form factors are experimentally unknown. Understanding such properties of the pentaquark baryons is, however, of great importance, since it will shed light on questions how the pentaquark baryons are constituted in terms of quarks and how they are produced in different reactions. In particular, the electric charge densities of the pentaquark baryons provide microscopical information on how their valence quarks and antiquark are located inside them, i.e., as already discussed in Ref. [17], electric properties of the pentaquark baryons reveal their internal quark structure.

Thus, in the present work, we aim at investigating the electric properties of the baryon anti-decuplet such as electric charge radii and electric form factors, based on the χ QSM with isospin symmetry and symmetry-conserving quantization [18] imposed. The χ QSM has been proved very successful not only in predicting the Θ^+ but also in describing various properties of SU(3) baryon octet and decuplet such as the mass splittings and form factors with the same set of fixed parameters [19]. In particular, the dependence of almost all

form factors on the momentum transfer is well reproduced within the χ QSM. As a result, the parity-violating asymmetries of polarized electron-proton scattering which require nine different form factors (six electromagnetic form factors $G_{E,M}^{(u,d,s)}(Q^2)$ and three axial-vector form factors $G_A^{(u,d,s)}(Q^2)$) are in good agreement with experimental data [20]. Therefore, it is worthwhile to extend the study of the form factors to the baryon anti-decuplet within the χ QSM. Moreover, as several models suggest, the pentaquark proton may contain a valence strange quark (s) and antiquark ($\bar{s}$), while the strange quark component of the proton is excited from the Dirac sea. Thus, it is also of great interest to study its strange electric form factor and to compare it with that of the proton.

The paper is organized as follows: In Section 2, we briefly review the formalism for the calculation of the electric properties of the baryon anti-decuplet in the χ QSM. In Section 3, we discuss the numerical results for the electric charge densities of the pentaquark baryons, and for their electric charge radii and electric form factors. We also compare the strange electric form factor of the pentaquark proton with that of the proton. In Section 4, we summarize and conclude the present work.

2. In this section we briefly review the formalism of the χ QSM for calculating the electric properties of the baryon anti-decuplet. We refer to Refs. [19, 21] for details. The electromagnetic form factors for the baryon anti-decuplet are expressed in terms of the quark matrix elements:

$$\langle B_{\overline{10}}(p') | J_\mu(0) | B_{\overline{10}}(p) \rangle = \bar{u}_{B_{\overline{10}}}(\mathbf{p}', s') \left[\gamma_\mu F_1^{B_{\overline{10}}}(Q^2) + i \frac{\sigma_{\mu\nu} q^\nu}{2M_N} F_2^{B_{\overline{10}}}(Q^2) \right] u_{B_{\overline{10}}}(\mathbf{p}, s), \quad (1)$$

where Q^2 is the square of the four momentum transfer $Q^2 = -q^2$ with $Q^2 > 0$. The M_N and $u_{B_{\overline{10}}}$ denote the nucleon mass and the Dirac spinor with the corresponding momentum $\mathbf{p}$ and spin s of the baryon anti-decuplet, respectively. The quark electromagnetic current J_μ in Euclidean space is defined in terms of the triplet ($J_\mu^{(3)}$) and octet ($J_\mu^{(8)}$) currents

$$J_\mu = -i\psi^\dagger \gamma_\mu \hat{Q} \psi = \frac{1}{2} \left(J_\mu^{(3)} + \frac{1}{\sqrt{3}} J_\mu^{(8)} \right) \quad (2)$$

with

$$J_\mu^{(3)} = -i\psi^\dagger \gamma_\mu \lambda_3 \psi, \quad J_\mu^{(8)} = -i\psi^\dagger \gamma_\mu \lambda_8 \psi. \quad (3)$$

The Dirac and Pauli form factors can be expressed in terms of the Sachs electric and magnetic form factors as follows:

$$\begin{aligned} G_E^{B_{\overline{10}}}(Q^2) &= F_1^{B_{\overline{10}}}(Q^2) - \frac{Q^2}{4M_N^2} F_2^{B_{\overline{10}}}(Q^2), \\ G_M^{B_{\overline{10}}}(Q^2) &= F_1^{B_{\overline{10}}}(Q^2) + F_2^{B_{\overline{10}}}(Q^2). \end{aligned} \quad (4)$$

The electric and magnetic form factors can be respectively written as matrix elements of the time and space components of the electromagnetic current:

$$\begin{aligned} \langle B_{\overline{10}}(p') | J_4(0) | B_{\overline{10}}(p) \rangle &= G_E^{B_{\overline{10}}}(Q^2) \delta_{s's} \\ \langle B_{\overline{10}}(p') | J_k(0) | B_{\overline{10}}(p) \rangle &= \frac{i}{2M_N} \epsilon_{klm} (\sigma^l)_{s's} q^m G_M^{B_{\overline{10}}}(Q^2), \end{aligned} \quad (5)$$

where σ^l stand for the Pauli spin matrices. Since we are interested in the electric form factors in this work, we concentrate on the first equation in Eq.(5).

Having performed a calculation as in Ref. [21], we arrive at the following expression:

$$G_E^{B_{\overline{10}}}(\mathbf{Q}^2) = G_E^{B_{\overline{10}},m_s^0}(\mathbf{Q}^2) + G_E^{B_{\overline{10}},m_s^1,\text{op}}(\mathbf{Q}^2) + G_E^{B_{\overline{10}},m_s^1,\text{wf}}(\mathbf{Q}^2), \quad (6)$$

where

$$G_E^{B_{\overline{10}},m_s^0}(\mathbf{Q}^2) = \frac{1}{8} \left(\frac{1}{3} \mathcal{B}(\mathbf{Q}^2) + \frac{\mathcal{I}_1(\mathbf{Q}^2)}{I_1} + 6 \frac{\mathcal{I}_2(\mathbf{Q}^2)}{I_2} \right) Q_{B_{\overline{10}}}, \quad (7)$$

$$\begin{aligned} G_E^{B_{\overline{10}},m_s^1,\text{op}}(\mathbf{Q}^2) = & \frac{m_s}{126I_1} \begin{bmatrix} 18 Q_{B_{\overline{10}}} \\ 11 Q_{B_{\overline{10}}} + 20 \\ 12 Q_{B_{\overline{10}}} + 32 \\ 13 Q_{B_{\overline{10}}} + 44 \end{bmatrix} (I_1 \mathcal{K}_1(\mathbf{Q}^2) - K_1 \mathcal{I}_1(\mathbf{Q}^2)) \\ & + \frac{m_s}{63I_2} \begin{bmatrix} 27 Q_{B_{\overline{10}}} \\ -Q_{B_{\overline{10}}} + 23 \\ -3 Q_{B_{\overline{10}}} + 20 \\ -5 Q_{B_{\overline{10}}} + 17 \end{bmatrix} (I_2 \mathcal{K}_2(\mathbf{Q}^2) - K_2 \mathcal{I}_2(\mathbf{Q}^2)) \\ & - m_s \left(\frac{1}{36} Q_{B_{\overline{10}}} - \frac{1}{252} \begin{bmatrix} 4 Q_{B_{\overline{10}}} \\ -3 Q_{B_{\overline{10}}} + 6 \\ -2 Q_{B_{\overline{10}}} + 4 \\ -Q_{B_{\overline{10}}} + 2 \end{bmatrix} \right) \mathcal{C}(\mathbf{Q}^2), \end{aligned} \quad (8)$$

$$\begin{aligned} G_E^{B_{\overline{10}},m_s^1,\text{wf}}(\mathbf{Q}^2) = & \frac{d_8}{6} \begin{bmatrix} 0 \\ -2 Q_{B_{\overline{10}}} + 2 \\ -Q_{B_{\overline{10}}} + 1 \\ 0 \end{bmatrix} \mathcal{B}(\mathbf{Q}^2) - \frac{d_8}{2} \begin{bmatrix} 0 \\ -2 Q_{B_{\overline{10}}} + 2 \\ -Q_{B_{\overline{10}}} + 1 \\ 0 \end{bmatrix} \frac{\mathcal{I}_1(\mathbf{Q}^2)}{I_1} \\ & - 2d_8 \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \frac{\mathcal{I}_2(\mathbf{Q}^2)}{I_2} + \frac{d_{27}}{120} \begin{bmatrix} 0 \\ 7 Q_{B_{\overline{10}}} - 2 \\ 6 Q_{B_{\overline{10}}} + 4 \\ 5 Q_{B_{\overline{10}}} + 10 \end{bmatrix} \mathcal{B}(\mathbf{Q}^2) \\ & - \frac{7d_{27}}{120} \begin{bmatrix} 0 \\ -7 Q_{B_{\overline{10}}} + 2 \\ -6 Q_{B_{\overline{10}}} - 4 \\ -5 Q_{B_{\overline{10}}} - 10 \end{bmatrix} \frac{\mathcal{I}_1(\mathbf{Q}^2)}{I_1} - \frac{d_{27}}{36} \begin{bmatrix} 0 \\ 21 Q_{B_{\overline{10}}} - 6 \\ 18 Q_{B_{\overline{10}}} + 12 \\ 15 Q_{B_{\overline{10}}} + 30 \end{bmatrix} \frac{\mathcal{I}_2(\mathbf{Q}^2)}{I_2} \\ & + \frac{d_{\overline{35}}}{112} \begin{bmatrix} 4 Q_{B_{\overline{10}}} \\ -3 Q_{B_{\overline{10}}} + 6 \\ -2 Q_{B_{\overline{10}}} + 4 \\ -Q_{B_{\overline{10}}} + 2 \end{bmatrix} \mathcal{B}(\mathbf{Q}^2) - \frac{d_{\overline{35}}}{112} \begin{bmatrix} 4 Q_{B_{\overline{10}}} \\ -3 Q_{B_{\overline{10}}} + 6 \\ -2 Q_{B_{\overline{10}}} + 4 \\ -1 Q_{B_{\overline{10}}} + 2 \end{bmatrix} \frac{\mathcal{I}_1(\mathbf{Q}^2)}{I_1} \\ & - \frac{d_{\overline{35}}}{56} \begin{bmatrix} 4 Q_{B_{\overline{10}}} \\ -3 Q_{B_{\overline{10}}} + 6 \\ -2 Q_{B_{\overline{10}}} + 4 \\ -1 Q_{B_{\overline{10}}} + 2 \end{bmatrix} \frac{\mathcal{I}_2(\mathbf{Q}^2)}{I_2} \end{aligned} \quad (9)$$

in the basis of $[\Theta^+, N_{\overline{10}}, \Sigma_{\overline{10}}, \Xi_{\overline{10}}]$. Here, the coefficients d_8 , d_{27} , and $d_{\overline{35}}$ are proportional to m_s and known from the SU(3) algebra [22], the I_1 etc. are moments of inertia whose expressions can be found in Ref [19], and the other quantities such as $\mathcal{B}(\mathbf{Q}^2)$ can be found in Ref. [23]. The $Q_{B_{\overline{10}}}$ denotes the charge of the corresponding pentaquark baryon.

3. We now discuss the results obtained from the present work. A detailed description of the numerical methods can be found in Ref. [19, 21]. We first want to emphasize that the only free parameter of the χ QSM is the constituent quark mass M . Since the electric charge radii and form factors turn out to be insensitive to the constituent quark mass M in general, we choose in this study the value $M = 420$ MeV, which is found from a best fit to many nucleon observables [19]. The other parameters of the model are the current nonstrange quark mass and the cut-off parameter of the proper-time regularization: They are all fixed for a given M in such a way that mesonic properties, i.e. the physical pion mass and decay constant, are well reproduced. The mass of the strange quark is throughout this work set to $m_s = 180$ MeV.




FIG. 1: Flavor-decomposed electric charge densities of the pentaquark baryon Θ^+ multiplied by r^2 as functions of r . The solid curves draw both the up (u) and down (d) quark components which are identical due to the assumed isospin symmetry, whereas the dashed curve depicts the strange (s) component (multiplied by (-1) for clarity). The vertical lines denote the position of the corresponding maximum values of the densities.

Figure 1 depicts the flavor-decomposed electric charge densities of the pentaquark baryon Θ^+ . The solid curves draw both the valence up ($\psi_u^\dagger\psi_u$) and down ($\psi_d^\dagger\psi_d$) quark densities which are identical due to the assumed isospin symmetry, whereas the dashed curve represents the strange quark ($\psi_s^\dagger\psi_s$) one. The vertical lines denote the corresponding maximum values of the densities. These flavor-decomposed densities of the Θ^+ make it possible for us to see the quark configuration inside the Θ^+ . As shown in Fig. 1, the peak of the valence strange anti-quark is located at slightly larger r , compared to that of the up and down quarks. It indicates that the strange anti-quark seems to be most probably found in the outer region of the Θ^+ , though it lies still rather close to the up and down quarks. Though we present here the densities of the Θ^+ , we find in general that the valence anti-quarks of the pentaquark baryons tend to be located at larger r and the shapes of the densities are very similar to those of the Θ^+ .

anti-decuplet	Θ^+	p_{10}^*	n_{10}^*	Σ_{10}^{*+}	Σ_{10}^{*0}	Σ_{10}^{*-}	Ξ_{10}^{*+}	Ξ_{10}^{*0}	Ξ_{10}^{*-}	Ξ_{10}^{*-}
$\langle r^2 \rangle_E [\text{fm}^2]$	0.770	0.771	0.014	0.772	0.008	0.756	0.773	0.003	0.768	0.769

TABLE I: The electric charge radii of the baryon anti-decuplet.

octet	p	n	Λ	Σ^+	Σ^0	Σ^-	Ξ^0	Ξ^-
$\langle r^2 \rangle_E [\text{fm}^2]$	0.768	-0.071	-0.029	0.771	0.026	0.720	-0.054	0.707

TABLE II: The electric charge radii of the baryon octet.

With these densities, we are able to calculate the electric charge radii $\langle r^2 \rangle_E$ of the baryon anti-decuplet which are listed in Table I. For comparison, we also list those of the baryon octet and decuplet in Table II and Table III, respectively. It is interesting to see that the electric charge radius of the Θ^+ is only slightly larger than that of the octet proton. This implies that the Θ^+ pentaquark baryon is quite compact, even though its main Fock state component corresponds to five quarks. In addition the SU(3) symmetry-breaking effects are negligible, as we will see later. We can also find from Table I that the electric charge radii of the baryon anti-decuplet are in general comparable to those of the baryon octet in magnitude apart from the neutral baryons. The electric charge radii of the neutral baryon anti-decuplet particles turn out to be much smaller than those of the neutral baryon octet ones. In particular, the $\langle r^2 \rangle_E^{\Xi_{10}^{*0}}$ is more than an order of magnitude smaller than $\langle r^2 \rangle_E^{\Xi^0}$. The reason lies in the fact that in the present formalism the electric charge radii of the neutral baryon anti-decuplet arise only from the SU(3) symmetry-breaking effects and are thus very tiny, in fact they vanish when the m_s corrections are switched off. We will discuss it more in detail later in the case of the electric form factors. The electric charge radii of the baryon decuplet are found to be larger than those of the octet and anti-decuplet.

Figure 2 draws the electric form factor of the Θ^+ with and without the current strange quark mass m_s . The effect of SU(3) symmetry breaking is almost negligible for the electric form factor of the Θ^+ . This feature holds basically for all electric form factors of the charged pentaquark baryons. The reason for this lies in the fact that in the χ QSM the strange quarks are not valence quarks but arise from sea quark excitations due to the rotations of the system.

In the left panel of Figure 3 we depict the electric form factors of the charged baryon anti-decuplet, while in its right panel we draw those of the neutral baryon anti-decuplet. It is interesting to see that there is almost no difference in the Q^2 dependence for charged pentaquark baryons with the same charge. It indicates that there is no flavor dependence. The reason can be found in Eqs.(7)-(9). The leading and rotational $1/N_c$ terms of the whole anti-decuplet are expressed in the first term of Eq.(7), which is just proportional to the charge of the corresponding pentaquark baryon $Q_{B_{10}}$. It means that the electric form factors of the pentaquark baryons with the same charge are identical when the SU(3) symmetry breaking is turned off. However, the effects of the SU(3) symmetry breaking on the electric form factors are in general small.

Moreover, the various terms of the m_s corrections are destructively interfering for the electric form factors of the baryon anti-decuplet as shown in Eqs.(8) and (9), so that the symmetry-breaking terms turn out to be negligibly small. As a result, the electric form factors of the charged pentaquark baryons appear to be almost the same.

In pure SU(3) symmetry, the electric form factors of the neutral pentaquark baryons vanish, since they are solely proportional to their charges. Thus, the electric form factor of

decuplet	Δ^{++}	Δ^+	Δ^0	Δ^-	Σ^{*+}	Σ^{*0}	Σ^{*-}	Ξ^{*0}	Ξ^{*-}	Ω^-
$\langle r^2 \rangle_E [\text{fm}^2]$	0.813	0.794	-0.038	0.869	0.823	-0.013	0.848	0.012	0.826	0.805

TABLE III: The electric charge radii of the baryon decuplet.




FIG. 2: Electric form factor of the Θ^+ with and without the strange quark mass. The solid curve depicts the form factor for $m_s = 180$ MeV, while the dashed one for $m_s = 0$.

a neutral pentaquark baryon play a role of measuring how much SU(3) symmetry is broken. As shown in the right panel of Fig. 3, the neutral electric form factors are negative, which leads to the positive electric charge radii (see Table I). We also find that the electric form factors of the neutral pentaquark baryons have almost the same Q^2 dependence.

Figure 4 compares the electric form factors of the nucleon with those of the pentaquark nucleon. The electric form factor of the pentaquark proton turns out to be almost the same as that of the proton. It is a remarkable result, reminding the fact that the pentaquark proton may contain the valence s and $\bar{s}$ quarks in addition to the valence uud component. We can infer from this result that the s and $\bar{s}$ overlap each other, so that they do not affect much the structure of the pentaquark proton. As a result, its general structure remains very similar to that of the proton.

In the right panel of Fig. 4, we compare the electric form factor of the pentaquark neutron to that of the octet neutron. As discussed above, the electric form factor of the anti-decuplet neutron vanishes in the SU(3) symmetric case, while that of the octet neutron does not. Though the general shapes of the electric form factors look similar, there is a large difference in the vicinity of $Q^2 = 0$. The neutron electric form factor changes sensitively near the zero momentum transfer as Q^2 increases, whereas that of the pentaquark neutron does slowly. It leads to the fact that the electric charge radius of the pentaquark neutron turns out to be very much smaller than that of the octet neutron.

Figure 5 compares the strange electric form factor of the pentaquark proton with that of the octet proton. Since in some models the pentaquark proton contains the valence s and $\bar{s}$




FIG. 3: Electric form factors of the baryon anti-decuplet as functions of Q^2 . In the left panel we depict those of the charged pentaquark baryons, while in the right panel we draw those of the neutral ones.




FIG. 4: Electric form factors of the nucleon and pentaquark nucleon as functions of Q^2 . In the left (right) panel we compare the proton (neutron) form factor with that of the pentaquark proton (neutron).

quarks, it may be expected that it has a larger strange component than the octet proton in which the strange quark contributions are bound to arise from the $s\bar{s}$ sea. In this context it is interesting that the strange electric form factor of the octet proton has been measured very recently [24]. As drawn in Fig. 5, the strange electric form factor of the pentaquark proton turns out to be larger than that of the proton by around 20 %, while its general shape is very similar to that of the proton. This is again caused by the fact that in the χ QSM the strange component is also a sea-quark excitation and not caused by a valence quark.

4. In the present work, we have applied the SU(3) chiral quark-soliton model (χ QSM) to




FIG. 5: Strange electric form factors of the proton and pentaquark proton as functions of Q^2 . The solid curve depicts that of the pentaquark proton, while the dashed one does that of the proton.

the calculation of electric form factors of the baryon anti-decuplet particles and their electric radii, emphasizing their difference from those of the baryon octet. We have assumed isospin asymmetry and have incorporated the symmetry-conserving quantization. The rotational $1/N_c$ and strange quark mass m_s corrections were taken into account. The dependence on the only free parameter M turned out very mild in general and we chose the proved value $M = 420$ MeV for which many properties of the octet and decuplet baryons are for many years known to be well reproduced.

We first have studied the flavor-decomposed electric density distributions of the pentaquark baryon Θ^+ , which provide information on the quark configuration inside the Θ^+ . We have found that the peak of the strange quark distribution is located 0.25 fm larger r , compared to that of the up and down quarks. It implies that the strange anti-quark is with higher probability found in the outer region of the Θ^+ , however still not too far from the up and down quarks.

We have calculated the electric charge radii of the baryon anti-decuplet. The results turned out that those of the charged anti-decuplet baryons are only slightly larger than those of the charged octet baryons. It means that the pentaquark baryons are quite compact objects, even though their leading Fock-component corresponds to five quarks. Apparently the higher Fock-components have a noticeable influence. Moreover, we found that the m_s corrections are almost negligible to the electric charge radii. This again indicates that the strange quarks arise from excitations of the Dirac sea rather than being valence objects. The electric charge radii of the neutral baryon anti-decuplet become much smaller than those of the neutral baryon octet. The reason can be found from the fact that the m_s corrections turn to be the leading-order contributions to them, i.e. when the m_s corrections are switched off, the electric charge radii of the neutral pentaquark baryons disappear. For comparison, we also presented the electric charge radii of the baryon decuplet, which are found to be

larger than those of the baryon octet and anti-decuplet.

We also have examined the electric form factors of the pentaquark baryons. Since the electric form factors of the anti-decuplet baryons are proportional to their charges in the SU(3) symmetric case, their general tendency is simply the same in SU(3) symmetry. Since the terms of the m_s corrections are destructively interfering, the SU(3) symmetry-breaking effects do not play any significant role in these form factors. However, for the neutral pentaquark baryons the m_s corrections turn out to provide the main contributions. Hence, these can be regarded as a measure of the degree of SU(3) symmetry breaking.

We have compared the electric form factors of the pentaquark nucleon with those of the nucleon. The electric form factor of the pentaquark proton are, remarkably, almost the same as that of the proton. It implies that the valence s and $\bar{s}$ quarks in the pentaquark proton overlap each other, so that they do not influence much the structure of the pentaquark proton. It means again that the pentaquark proton is a compact baryon despite of its five-quark structure. As for the neutron electric form factors, their general shapes look similar. However, it is found that there is a large difference in the vicinity of $Q^2 = 0$ where the form factor of the octet neutron changes much faster than the one of the anti-decuplet neutron.

We finally have investigated the strange electric form factors of the pentaquark proton. The pentaquark proton may contain the valence s and $\bar{s}$ quarks, whereas the strange quark component of the proton arises from the $s\bar{s}$ sea. Thus, we anticipate that the pentaquark proton might have a larger strange component than that of the proton. Indeed, the strange electric form factor of the pentaquark proton turns out to be larger than that of the proton by around 20 %.

In conclusion, we have found two significant features of the pentaquark baryons by studying their electric properties: Firstly, even though their leading Fock-component consists of five quarks, they are rather compact baryons, being comparable to the baryon octet. Secondly, the effects of SU(3) symmetry breaking do not play any important role in the case of the charged pentaquark baryons. However, those effects are leading-order contributions for the neutral ones. As a result, electric properties of the neutral pentaquark baryons show the extent of SU(3) symmetry breaking. We plan further studies on other properties of the pentaquark baryons such as magnetic transition and axial-vector form factors, which will appear soon elsewhere.

Acknowledgments

The authors are very grateful to M. Praszalowicz for helpful discussions and comments. The present work is supported by a Korean-German (F01-2004-000-00102-0) grant of the KOSEF and of the Deutsche Forschungsgemeinschaft (DFG). The work of HChK is also supported by the Korea Research Foundation Grant funded by the Korean Government(MOEHRD) (KRF-2005-202-C00102). The work is partly supported by the Transregio-Sonderforschungsbereich Bonn-Bochum-Giessen, the Verbundforschung of the Federal Ministry for Education and Research (BMBF) of Germany, the Graduiertenkolleg Bochum-Dortmund, the COSY-project Jülich as well as the EU Integrated Infrastructure Initiative

- [1] D. Diakonov, V. Petrov, and M. V. Polyakov, Z. Phys. A **359**, 305 (1997).
- [2] M. Praszalowicz, in: Workshop on Skyrmions and Anomalies, M. Jezabek and M. Praszalowicz (Eds.), World Scientific, 1987, p.112. Phys. Lett. B **575**, 234 (2003).
- [3] K. H. Hicks, Prog. Part. Nucl. Phys. **55**, 647 (2005) and references therein.
- [4] T. Nakano *et al.* [LEPS Collaboration], Phys. Rev. Lett. **91**, 012002 (2003).
- [5] C. Alt *et al.* [NA49 Collaboration], Phys. Rev. Lett. **92**, 042003 (2004).
- [6] M. Battaglieri *et al.* [CLAS Collaboration], arXiv:hep-ex/0510061.
- [7] T. Hotta [LEPS Collaboration], Acta Phys. Polon. B **36**, 2173 (2005).
- [8] K. Miwa *et al.* [KEK-PS E522 Collaboration], arXiv:nucl-ex/0601032.
- [9] V. Kuznetsov [GRAAL Collaboration], arXiv:hep-ex/0409032.
- [10] V. Kuznetsov *et al.*, arXiv:hep-ex/0601002.
- [11] H.-Ch. Kim, M. Polyakov, M. Praszalowicz, G. S. Yang and K. Goeke, Phys. Rev. D **71**, 094023 (2005).
- [12] Y. Azimov, V. Kuznetsov, M. V. Polyakov and I. Strakovsky, Eur. Phys. J. A **25**, 325 (2005).
- [13] K. S. Choi, S. i. Nam, A. Hosaka and H.-Ch. Kim, arXiv:hep-ph/0512136, to appear in Phys. Lett. B.
- [14] B. K. Jennings and K. Maltman, Phys. Rev. D **69**, 094020 (2004) and references therein.
- [15] S. L. Zhu, Int. J. Mod. Phys. A **19**, 3439 (2004).
- [16] K. Goeke, H.-Ch. Kim, M. Praszalowicz and G. S. Yang, Prog. Part. Nucl. Phys. **55**, 350 (2005), and references therein.
- [17] M. V. Polyakov, arXiv:hep-ph/0412274.
- [18] M. Praszalowicz, T. Watabe and K. Goeke, Nucl. Phys. A **647**, 49 (1999).
- [19] C. V. Christov *et al.*, Prog. Part. Nucl. Phys. **37**, 91 (1996).
- [20] A. Silva, H.-Ch. Kim, D. Urbano and K. Goeke, arXiv:hep-ph/0601239.
- [21] H.-Ch. Kim, A. Blotz, M. V. Polyakov and K. Goeke, Phys. Rev. D **53**, 4013 (1996).
- [22] G. S. Yang, H.-Ch. Kim, M. Praszalowicz and K. Goeke, Phys. Rev. D **70**, 114002 (2004).
- [23] A. Silva, H.-Ch. Kim and K. Goeke, Eur. Phys. J. A **22**, 481 (2004).
- [24] K. A. Aniol *et al.* [HAPPEX Collaboration], arXiv:nucl-ex/0506010.